

## Wessex Regional Genetics Laboratory Publications 2018

**Bateman MS, Collinson MN, Bunyan DJ, Collins AL, Duncan P, Firth R, Harrison V, Homfrey T, Huang S, Kirk B, Lachlan KL, Maloney VK, Barber JCK.** Incomplete penetrance, variable expressivity, or dosage insensitivity in four families with directly transmitted unbalanced chromosome abnormalities. *Am J Med Genet Part A* 2018; **176A**: 319-329

Begemann M, Rezwan FI, Beygo J, Docherty LE, Kolarova J, Schroeder C, Buiting K, Chokkalingam K, Degenhardt F, Wakeling EL, Kleinle S, Fassrainer *Med Otolaryngol Head Neck Surg* 2018; **55**: 497-504

Brioude F, Hennekam R, Blieff J, Coze C, Eggermann T, Ferrero GB, Kratz C, LeBouc Y, Maas SM, Mackay DJG, Maher ER, Mussa A, Netchine I. Revisiting Wilms tumour surveillance in Beckwith-Wiedemann syndrome with IC2 methylation loss. *Eur J Clin Endocrinol* 2018; **146**: 471-471

Brioude F, Kalish JM, Mussa A, Foster AC, Bliek J, Ferrero GB, Boonen SE, Cole T, Baker R, Bertoletti M, Cocchi G, Coze C, De Pellegrin M, Hussain K, Ibrahim A, Kilby MD, Krajewska-Walasek M, Kratz CP, Ladusans EJ, Lapunzina P, Le Bouc Y, Maas SM, Macdonald F, Önup K, Peruzzi L, Rossignol S, Russo S, Shipster C, Skórka A, Tatton-Brown K, Tenorio J, Tortora C, Grönakoc K, Netchine I, Hennekam RC, Prawitt D, Tümer Z, Eggermann T, Mackay DJG, Riccio A, Maher ER. Clinical and molecular diagnosis, screening and management of Beckwith-Wiedemann syndrome: an international consensus statement. *Nat Rev Endocrinol* 2018; **16**: 166 doi:10.1038/nrendo.2017.166

Catherwood MA, Graham A, Cuthbert RJG, Garrec C, Gardie B, Girodon F, Laird S, Cross NCP, McMullin MF. Absence of CALR mutations in idiopathic erythrocytosis patients with low serum erythropoietin levels. *Acta Endocrinol* 2018; **139**: 217-219

Ceroni F, Aguilera-Garcia D, Chassaing N, Bax DA, Blanco-Kelly F, Ramnos P, Tarilonte M, Villaverde C, da Silva LRJ, Ballesta-Martinez MJ, Sanchez-Soler MJ, Holt RJ, Cooper-Charles L, Bruty J, Wallis Y, McMullan D, Hoffman J, Bunyan D, Stewart A, Stewart H, Lachlan K, Study DDD, Fryer A, McKay V, Roume J, Dureau P, Saggar A, Griffiths M, Calvas P, Ayuso C, Corton M, Ragge NK. New *A* variants and phenotypes highlight its critical role in a broad spectrum of eye anomalies. 2018 doi:org/10.1007/s00439-018-1875-2.

Cross NCP, White HE, Evans PAS, Hancock J, Copland M, Milojkovic D, Mason J, Craine S, Mead AJ. Consensus on BC-ABL reporting in chronic myeloid leukaemia in the UK. *Br J Haematol* 2018; **173**: 15542 doi:10.1111/bjh.15542

Gast C, Marinaki A, Arenas-Hernandez M, Campbell S, Seaby EG, Pengelly RJ, Gale DP, Connor TM, Bunyan DJ, Hoda

ová K, Živná M, Kmoch S, Ennis S, Venkat-Raman G. Autosomal dominant tubulointerstitial kidney disease-UMOD is the most frequent non-polycystic genetic kidney disease. *Nat Rev Endocrinol* 2018 doi:org/1.1186/s12882-018-1107-y

Hoade Y, Metzgeroth G, Schwaab J, Reiter A, Cross NCP. Routine screening for M541L is not warranted in the diagnostic work-up of patients with hypereosinophilia. 2018 *Acta Endocrinol* 2018; **139**: 71-73

Khodeer A, Al Obaidi M, Babb A, Liu C, Cross NCP, Bain BJ. Chronic neutrophilic leumemia  
A  e o 2018 **93**: 841-842

Koczkowska M, Chen Y, Callens T, Gomes A, Sharp A, Johnson S, Hsiao M-C, Chen Z,  
Balasubramanian M, Barnett CP, Becker TA, Ben-Shachar S, Be19.7187(h)